

# Biostatistics 615/815 Lecture 18:

## Single Dimensional and Multidimensional Optimizations

Hyun Min Kang

March 21th, 2011

Hyun Min Kang

Biostatistics 615/815 - Lecture 18

March 21th, 2011 1 / 39

## Recap : Root Finding with C++

```
double binaryZero(myFunc foo, double lo, double hi, double e) {
    for (int i=0;; ++i) {
        double d = hi - lo; // f(lo) < 0, f(hi) > 0, d can be positive or negative
        double point = lo + d * 0.5; // d is + for increasing func, - for decreasing
        double fpoint = foo(point); // evaluate the value of the function
        if (fpoint < 0.0) {
            d = lo - point; lo = point; //
        }
        else {
            d = point - hi; hi = point;
        }
        // e is tolerance level (higher e makes it faster but less accurate)
        if (fabs(d) < e || fpoint == 0.0) {
            std::cout << "Iteration " << i << ", point = " << point
            << ", d = " << d << std::endl;
            return point;
        }
    }
}
```

Hyun Min Kang

Biostatistics 615/815 - Lecture 18

March 21th, 2011 3 / 39

Hyun Min Kang

Biostatistics 615/815 - Lecture 18

March 21th, 2011 2 / 39

## Recap : Improvements to Root Finding

### Approximation using linear interpolation

$$f^*(x) = f(a) + (x - a) \frac{f(b) - f(a)}{b - a}$$

### Root Finding Strategy

- Select a new trial point such that  $f^*(x) = 0$

Hyun Min Kang

Biostatistics 615/815 - Lecture 18

March 21th, 2011 4 / 39

## Annoucements

### Homework and Grading

- Homework #5 announced
- Homework grading is still pending

### Thursday March 24th

- Mary Kate Trost will introduce us a very useful C++ library


## Recap : Detailed Minimization Strategy

- ① Find 3 points such that
  - $a < b < c$
  - $f(b) < f(a)$  and  $f(b) < f(c)$
- ② Then search for minimum by
  - Selecting trial point in the interval
  - Keep minimum and flanking points

## Recap : Golden Search

```
double goldenSearch(myFunc foo, double a, double b, double c, double e) {
    int i = 0;
    double fb = foo(b);
    while ( fabs(c-a) > fabs(b*e) ) {
        double x = b + goldenStep(a, b, c);
        double fx = foo(x);
        if ( fx < fb ) {
            (x > b) ? ( a = b ) : ( c = b );
            b = x; fb = fx;
        }
        else {
            (x < b) ? ( a = x ) : ( c = x );
        }
        ++i;
    }
    std::cout << "i = " << i << ", b = " << b << ", f(b) = " << foo(b) << std::endl;
    return b;
}
```

## Recap : The Golden Search



## Today

### A better single-dimensional optimization

- Parabolic interpolation
- Adaptive method

### Multi-dimensional optimization

- Simplex algorithm

## Better optimization using local approximation

- Root finding example
  - Binary search reduces the search space by constant factor 1/2
  - Linear approximation may reduce the search space more rapidly for most well-defined functions
- Minimization problem
  - Golden search reduces the search space by 38%
  - Using a quadratic approximation of the function may achieve better optimization results

## Parabolic Approximation


$$f^*(x) = Ax + Bx + C$$

The value minimizes  $f^*(x)$  is

$$x_{min} = -\frac{B}{2A}$$

This strategy is called "inverse parabolic interpolation"

## Approximation using parabola



## Fitting a parabola

- Can be fitted with three points
- Points must not be co-linear
- $f^*(x_1) = f(x_1), f^*(x_2) = f(x_2), f^*(x_3) = f(x_3)$ .

$$\begin{aligned} C &= f(x_1) - Ax_1^2 - Bx_1 \\ B &= \frac{A(x_2^2 - x_1^2) + f(x_1) - f(x_2)}{x_1 - x_2} \\ A &= \frac{f(x_3) - f(x_2)}{(x_3 - x_2)(x_3 - x_1)} - \frac{f(x_1) - f(x_2)}{(x_1 - x_2)(x_3 - x_1)} \end{aligned}$$

## Minimum for a Parabola

- General expression for finding minimum of a parabola fitted through three points

$$x_{min} = x_2 - \frac{1}{2} \frac{(x_2 - x_1)^2(f(x_2) - f(x_1)) - (x_2 - x_3)^2(f(x_2) - f(x_1))}{(x_2 - x_1)(f(x_2) - f(x_1)) - (x_2 - x_3)(f(x_2) - f(x_1))}$$

## Avoiding degenerate case

- Fitted minimum could overlap with one of original points
- Ensure that each new point is distinct from previously examined points

## Fitting a Parabola

```
// Returns the distance between b and the abscissa for the
// fitted minimum using parabolic interpolation
double parabola_step (double a, double fa, double b, double fb, double c, double f)
{
    // Quantities for placing minimum of fitted parabola
    double p = (b - a) * (fb - fc);
    double q = (b - c) * (fb - fa);
    double x = (b - c) * q - (b - a) * p;
    double y = 2.0 * (p - q);
    // Check that q is not zero
    if (fabs(y) < ZEPS)
        return golden_step (a, b, c);
    else
        return x / y;
}
```

## Avoiding degenerate steps

```
double adjust_step(double a, double b, double c, double step, double e) {
    double min_step = fabs(e * b) + ZEPS;
    if (fabs(step) < min_step)
        return step > 0 ? min_step : -min_step;
    // If the step ends up too close to previous points,
    // return zero to force a golden ratio step ...
    if (fabs(b + step - a) <= e || fabs(b + step - c) <= e)
        return 0.0;
    return step;
}
```

## Generating New Points

- Use parabolic interpolation by default
- Check whether improvement is slow
- If step sizes are not decreasing rapidly enough, switch to golden section

## Overall

The main function simply has to

- Generate new points using building blocks
- Update the triplet bracketing the minimum
- Check for convergence

## Adaptive calculation of step size

```
double calculate_step(double a, double fa, double b, double fb,
                      double c, double fc, double last_step, double e) {
    double step = parabola_step(a, fa, b, fb, c, fc);
    step = adjust_step(a, b, c, step, e);
    if (fabs(step) > fabs(0.5 * last_step) || step == 0.0)
        step = golden_step(a, b, c);
    return step;
}
```

## Overall Minimization Routine

```
double find_minimum(myFunc foo, double a, double b, double c, double e) {
    double fa = foo(a), fb = foo(b), fc = foo(c);
    double step1 = (c - a) * 0.5, step2 = (c - a) * 0.5;
    while ( fabs(c - a) > fabs(b * e) + ZEPS) {
        double step = calculate_step (a, fa, b, fb, c, fc, step2, e);
        double x = b + step;
        double fx = foo(x);
        if (fx < fb) {
            if (x > b) { a = b; fa = fb; }
            else { c = b; fc = fb; }
            b = x; fb = fx;
        }
        else {
            if (x < b) { a = x; fa = fx; }
            else { c = x; fc = fx; }
            step2 = step1; step1 = step;
        }
    }
    return b; }
```

## Important Characteristics


- Parabolic interpolation often converges faster
  - The preferred algorithm
- Golden search provides worst-case performance guarantee
  - A fall-back for uncooperative functions
- Switch algorithms when convergence is slow
- Avoid testing points that are too close

## Recommended Reading

- Numerical Recipes in C++
- Chapter 10.0 - 10.3

## More advanced strategy : Brent's algorithm

- Track 6 points (not all distinct)
  - The bracket boundaries ( $a, b$ )
  - The current minimum  $x$
  - The second and third smallest value ( $w, v$ )
  - The new points to be examined  $u$
- Parabolic interpolation
  - Using  $(x, w, v)$  to propose new value for  $u$ .
  - Additional care is required to ensure  $u$  falls between  $a$  and  $b$ .



## A general mixture distribution

$$p(x; \pi, \phi, \eta) = \sum_{i=1}^k \pi_i f(x; \phi_i, \eta)$$

*x* observed data

$\pi$  mixture proportion of each component

*f* the probability density function

$\phi$  parameters specific to each component

$\eta$  parameters shared among components

*k* number of mixture components

## Problem : Maximum Likelihood Estimation

### Finding Maximum-likelihood

Find parameters that maximizes the likelihood of the entire sample

$$L = \prod_i p(x_i | \pi, \phi, \eta)$$

### Calculating in log-space

Or equivalently, consider log-likelihood to avoid underflow

$$l = \sum_i \log p(x_i | \pi, \phi, \eta)$$

## Gaussian MLE in single-dimensional space

$$p(x; \mu, \sigma^2) = \mathcal{N}(x; \mu, \sigma^2)$$

Given  $x$ , what is the MLE parameters of  $\mu$  and  $\sigma^2$ ?

- Analytical solution does exist
- $\hat{\mu} = \sum_{i=1}^n x_i / n$
- $\hat{\sigma}^2 = \sum_{i=1}^n (x_i - \hat{\mu})^2 / n$

## MLE in Gaussian mixture

### Parameter estimation in Gaussian mixture

- No analytical solution
- Numerical optimization required
- Multi-dimensional optimization problem
  - $\mu_1, \mu_2, \dots, \mu_k$
  - $\sigma_1^2, \sigma_2^2, \dots, \sigma_k^2$


### Possible approaches

- Simplex Method
- Expectation Maximization
- Markov-Chain Monte Carlo

## The Simplex Method

- Calculate likelihoods at simplex vertices
  - Geometric shape with  $k+1$  corners
  - A triangle in  $k=2$  dimensions
- Simplex *crawls*
  - Towards minimum
  - Away from maximum
- Probably the most widely used optimization method


## How the Simplex Method Works




## Simplex Method in Two Dimensions

- Evaluate functions at three vertices
  - The highest (worst) point
  - The next highest point
  - The lowest (best) point
- Intuition
  - Move away from high point, towards low point


## Direction for Optimization




## Reflection




## Reflection and Expansion



## Contaction (1-dimension)




## Contaction



If a simple contraction doesn't improve things, then try moving all points towards the current minimum

## Summary : The Simplex Method



## Today

- Single-dimensional minimization
  - Minimization using Parabola
  - Adaptive minimization using parabola and golden search
  - A taste of Brent's method
- Multi-dimensional optimization
  - Gaussian mixture example
  - Simplex algorithm

## Upcoming lectures

### Next Lecture

- Special lecturer : Mary Kate Trost
- Practical lessons in C++

### Next week

- Details of simplex algorithm
- Expectation-Maximization algorithm